

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281108132>

# First report of Tomato leaf curl Sinaloa virus infecting tomato crops in Panama

Article · June 2015

DOI: 10.5197/j.2044-0588.2015.031.030

CITATIONS

8

READS

135

7 authors, including:



**Jose Angel Herrera Vasquez**

Instituto de Investigación Agropecuaria de Panamá

35 PUBLICATIONS 259 CITATIONS

[SEE PROFILE](#)



**Deibis Yorlenis Ortega Castro**

Universitat Politècnica de València

3 PUBLICATIONS 20 CITATIONS

[SEE PROFILE](#)



**Salvatore Walter Davino**

Università degli Studi di Palermo

123 PUBLICATIONS 781 CITATIONS

[SEE PROFILE](#)



**Luis C Mejía**

Instituto de Investigaciones Científicas y Servicios de Alta Tecnología

52 PUBLICATIONS 2,204 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



predoctoral project [View project](#)



Influence of landscape, climatology and agricultural management on biodiversity linked to vineyards: effects on ecosystem services and sustainability under Global Change [View project](#)



# First report of *Tomato leaf curl Sinaloa virus* infecting tomato crops in Panama

J.A. Herrera-Vásquez<sup>1\*</sup>, D. Ortega<sup>2</sup>, A.B. Romero<sup>1</sup>, S. Davino<sup>3,4</sup>, L.C. Mejía<sup>5,6</sup>, S. Panno<sup>4</sup> and M. Davino<sup>7</sup>

<sup>1</sup> Laboratorio de Protección Vegetal (LPV), Centro de Investigación Agropecuaria Central (CIAC), Instituto de Investigación Agropecuaria de Panamá (IDIAP), Ctra. Panamericana, Los Canelos, Santa María, Estafeta de Divisa, 0619 Herrera, Panama; <sup>2</sup> Escuela de Biología, Facultad de Ciencias Naturales, Exactas y Tecnología, Universidad de Panamá (UP), Ciudad Universitaria Dr. Octavio Méndez Pereira, Estafeta Universitaria, 3366 Panama 4, Panama; <sup>3</sup> Department of Agricultural and Forestry Science, University of Palermo, Viale delle Scienze Ed 5, 90128 Palermo, Italy; <sup>4</sup> Euro-Mediterranean Institute of Science and Technology, Via Emerico Amari 123, 90139 Palermo, Italy; <sup>5</sup> Centro de Biología Molecular y Celular de Enfermedades, Instituto de Investigaciones Científicas y Servicios de Alta Tecnología (INDICASAT AIP), Edificio 219, Ciudad del Saber, Clayton, PO Box 0843-01103 Panama 5, Panama; <sup>6</sup> Smithsonian Tropical Research Institute (STRI), Balboa, Ancon, PO Box 0843-03092 Panama, Panama; <sup>7</sup> Dipartimento DISPA, University of Catania, Via S. Sofia 100, 95123 Catania, Italy

\*E-mail: joshervs11@gmail.com

Received: 31 Dec 2014. Published: 21 Jun 2015. Keywords: *Solanum lycopersicum*, viral plant disease

In April 2011 and September 2012, virus-like symptoms were observed in open field- and greenhouse-grown tomato crops (*Solanum lycopersicum*) in Chiriquí, the westernmost province of Panama. Samples from symptom-bearing plants (127 in all) were collected and tested for the presence of begomoviruses by polymerase chain reaction (PCR) assays with sets of degenerated primers designed to amplify parts of the DNA-A and DNA-B components (Rojas *et al.*, 1993; Table 1). Products of the expected sizes, obtained with both DNA-A- and DNA-B-specific primers for 49 samples, suggested infection with New World bipartite begomoviruses. This corresponds to an incidence of 26% (8 plants) in open field, and 43% (41 plants) in greenhouse crops. Primers specific for ten tomato-infecting begomoviruses found in Central America (Engel *et al.*, 1998; Nakhla *et al.*, 2005; Table 1) were used to typify the PCR-positive samples.

This analysis revealed *Potato yellow mosaic Panama virus* (PYMPV) or *Tomato leaf curl Sinaloa virus* (ToLCSiV) in 44 (90%) or 40 (82%) of the samples, respectively. All contained at least one virus, the majority (i.e. 35) indeed both, with no indication of the other viruses tested. BLAST analysis of two PCR products' sequences of the distinct viruses (GenBank Accession Nos. KP313717 for PYMPV and KP318651 for ToLCSiV, respectively) revealed that KP313717 shared 99% DNA sequence identity with PYMPV - [Panama:Divisa:Tomato:1996] (PYMPV-[PA:Div:Tom:96], (Y15034) (Engel *et al.*, 1998), and KP318651 99% identity with ToLCSiV - [Nicaragua:Santa Lucia] (ToLCSiV-[NI:SL], (AJ608286) (Rojas *et al.*, 2005) and three other ToLCSiV sequences: [Nicaragua:Santa Lucia] (AJ508779), [Nicaragua:Sebaco] (AJ508780) (Rojas *et al.*, 2005), and [Costa Rica:Alajuela] (AF131213); as well as 98% identity with ToLCSiV-[Nicaragua:Condega] (AJ508778) (Rojas *et al.*, 2005).

Differences in symptom expression were in some cases observed between plants infected with both viruses (Fig. 1A), or with PYMPV (Fig. 1B) or ToLCSiV (Fig. 1C) alone. Begomovirus-free plants (Fig. 1D) also showed virus-like symptoms resembling those induced by other viruses, especially in the *Potyviridae* and *Tobamoviridae* (Polston & Anderson, 1997). Due to the high capacity of recombination between different begomoviruses (Davino *et al.*, 2012), the existence or development of novel recombinant molecules cannot be excluded, which could lead to the emergence of new begomoviruses with different biological properties compared to the

ancestral parental viruses in the future. To our knowledge, this is not only the first detection of ToLCSiV in Panama, but also the first report of PYMPV in Panama's western highlands, and the first ever report of PYMPV/ToLCSiV mixed infection. Additional studies on incidence and distribution of these viruses in Panama are in progress.

## Acknowledgements

This study was funded by FONTAGRO and SENACYT (Panama). J.A.H.-V. and L.C.M. were sponsored by the SENACYT SNI program.

## References

- Davino S, Miozzi L, Panno S, Rubio L, Davino M, Accotto GP, 2012. Recombination profiles between *Tomato yellow leaf curl virus* and *Tomato yellow leaf curl Sardinia virus* in laboratory and field conditions: evolutionary and taxonomic implications. *Journal of General Virology* **93**, 2712-2717. <http://dx.doi.org/10.1099/vir.0.045773-0>
- Engel M, Fernández O, Jeske H, Frischmuth T, 1998. Molecular characterization of a new whitefly-transmissible bipartite geminivirus infecting tomato in Panama. *Journal of General Virology* **79**, 2313-2317.
- Nakhla MK, Sorenson A, Mejía L, Ramírez P, Karkashian JP, Maxwell DP, 2005. Molecular characterization of tomato-infecting begomoviruses in Central America and development of DNA-based detection methods. *Acta Horticulturae* **695**, 277-288.
- Polston JE, Anderson PK, 1997. The emergence of whitefly-transmitted geminiviruses in tomato in the Western Hemisphere. *Plant Disease* **81**, 1358-1369. <http://dx.doi.org/10.1094/PDIS.1997.81.12.1358>
- Rojas MR, Gilbertson RL, Russell DR, Maxwell DP, 1993. Use of degenerate primers in the polymerase chain reaction to detect whitefly-transmitted geminiviruses. *Plant Disease* **77**, 340-347. <http://dx.doi.org/10.1094/PD-77-0340>
- Rojas A, Kvarnheden A, Marcenaro D, Valkonen JP, 2005. Sequence characterization of Tomato leaf curl Sinaloa virus and *Tomato severe leaf curl virus*: Phylogeny of New World begomoviruses and detection of recombination. *Archives of Virology* **150**, 1281-1299. <http://dx.doi.org/10.1007/s00705-005-0509-x>



Figure 1

**To cite this report:** Herrera-Vásquez JA, Ortega D, Romero AB, Davino S, Mejía LC, Panno S, Davino M, 2015. First report of *Tomato leaf curl Sinaloa virus* infecting tomato crops in Panama. *New Disease Reports* **31**, 30. <http://dx.doi.org/10.5197/j.2044-0588.2015.031.030>  
©2015 The Authors

This report was published on-line at [www.ndrs.org.uk](http://www.ndrs.org.uk) where high quality versions of the figures can be found.